

IDENTIFIED PARTICLE PRODUCTION AT HIGH TRANSVERSE MOMENTUM IN NUCLEUS-NUCLEUS COLLISIONS AT RHIC

V.S. Pantuev

*Department of Physics and Astronomy, Stony Brook University,
Stony Brook, New York, 11794-3800, USA*

Experimental results for identified hadron spectra in AA collisions at RHIC are reviewed. Jet suppression in central AuAu collisions at 200 GeV is seen in leading meson and baryon spectra at high p_t . Enhanced baryon production at intermediate p_T ($2 < p_t < 5$ GeV/c) is related to quark content but not its mass. In d+Au collisions Cronin enhancement is observed with larger magnitude for baryons than for mesons, but this difference can not explain the baryon-to-meson ratio in Au+Au collisions. In Au+Au the observations at 62.4 GeV and 200 GeV are very similar, but smaller suppression of the leading particles is seen at low beam energy, which is evidence for a smooth beam energy dependence of the effect. Near and away side jets seen in two-particle correlations are very similar for the leading baryon and mesons.

Since the year 2000 four experiments at the Relativistic Heavy Ion Collider (RHIC) at Brookhaven National Laboratory are searching for a new state of nuclear matter: the quark-gluon plasma. All experiments confirmed the effect of particle suppression at high transverse momenta in the most central Au+Au collisions¹. This phenomenon is consistent with the general expectation from pQCD calculations for final-state partonic energy loss or “jet quenching” in produced dense matter. At the same time it was found that protons and antiprotons are not suppressed as pions², Figure 1. Here the parameter R_{cp} is defined as:

$$R_{cp}(p_T) = \frac{(1/\langle N_{binary}^{cent} \rangle) d^2 N_{cent}^{A+A} / dp_T d\eta}{(1/\langle N_{binary}^{periph} \rangle) d^2 N_{periph}^{A+A} / dp_T d\eta}. \quad (1)$$

Jet production, and hence its leading particle, has a small cross section and should be proportional to the number of binary nucleon-nucleon collisions, N_{binary} , for each AA collision centrality. Thus, to see the nuclear modification effect versus centrality, particle yields in R_{cp} are normalized to N_{binary} .

To understand whether the proton non-suppression is a mass effect due to an additional radial flow or a pure quark-content effect (baryon vs. meson), many measurements have been done with other particles^{3,4}, Figure 2 and Figure 3. First, data confirm that this is a baryon effect, probably related to the number of constituent quarks, three for baryons and two for mesons. Second, baryon suppression is also observed, but it starts at higher momenta above 4 GeV/c. The baryon enhancement seen in Au+Au at 2-4 GeV/c is a feature of the transition region between soft and hard scattering processes.

To confirm that jet quenching is indeed a final state effect in central Au+Au collisions, a set of “control” experiments at RHIC has been performed with d+Au beams. If the suppression is an initial state effect, then it should be seen in low density d+Au collisions also. At midrapidity




Figure 1: Central to peripheral yield ratio per binary collision, R_{CP} , vs. transverse momentum for neutral pions and protons in Au+Au collisions at 200 GeV.




Figure 2: Central to peripheral yield ratio per collision vs. transverse momentum for charge mesons and baryons in Au+Au collisions at 200 GeV.

no suppression was observed. Even more, some enhancement was seen for more central d+Au events⁵, Figure 4. This, so called, Cronin effect was observed already in mid 1970-th in proton beams on nuclear targets. The enhancement is larger for baryons than for mesons but this is a small effect and can not explain the factor 4-5 baryon-meson difference at intermediate p_t in central Au+Au collisions.

The observation of strong meson suppression at high transverse momenta in Au+Au collisions at 200 GeV gave rise to the question about the beam energy dependance of a such effect. RHIC performed a short one week Au+Au run at $\sqrt{s} = 62.4$ GeV. Preliminary results, Figure 5, show a significant suppression of the neutral pion yield for the most central collisions⁶, but the suppression factor at 62.4 GeV is less than at 200 GeV. This is evidence for a smooth beam energy dependence of the effect. At the same time, at 62.4 GeV the nuclear modification factor for charged hadrons, i.e. pions, kaons and protons, does not show suppression in the measured range of $p_t < 4$ GeV/c. An explanation of this feature is presented in Figure 6: in most central events at high momentum the charged hadron yield contains more protons than pions. There is no direct measurements of the nuclear modification factor for identified baryons at $\sqrt{s} = 62.4$ GeV, but from Figure 5 and Figure 6 one may conclude, that baryon production, or at least for protons, is similar at 62.4 and 200 GeV.

The very large particle multiplicity in Au+Au collisions at RHIC does not allow for a full jet reconstruction - the background of soft particles is too high. Nevertheless, it was demonstrated that two particle azimuthal correlation for large transverse momentum particles could be a valid tool for jet identification⁷. 2-particle azimuthal correlations allow to investigate the centrality dependence, the shape and yield in forward and backward jets, to measure the spectrum of particles in the jet. The results for the particle yield in the jets^{8,9} are presented in Figure 7. The leading particle was identified as a meson or baryon, the second particle was a non-identified charged hadron. The particle yield per trigger in the near side and away side jets is almost the same for identified leading mesons and baryons, at least in the mesured momentum range, regardless of a large difference in the inclusive baryon and meson yields at $2 < p_t < 5$ GeV/c. Different versions of the recombination/coalescence parton model were proposed to describe the enhanced p/π ratio for inclusive spectra¹⁰. This model goes beyond the standard picture of high transverse momentum hadron production by fragmentation of the energetic parton. It proposes that hadron production at a few GeV/c in an enviroment with high density of partons




Figure 3: Central to peripheral yield ratio per collision vs. transverse momentum for pions, protons and ϕ -mesons in Au+Au collisions at 200 GeV.




Figure 4: Central to peripheral yield ratio per binary collision vs. transverse momentum for hyperons, kaons and ϕ -mesons in d+Au collisions at 200 GeV. Dotted line at $R_{CP} = 1$ shows expectation from binary scaling.

occurs via the parton recombination. Another semi-phenomenological model describes the large hadron/pion ratio by a non-universal parton transverse momentum broadening¹¹. Authors of this paper argue that smearing of the parton intrinsic k_T in pA and AA collisions should be larger for protons than for pions. Less schematic calculations should be done to test both models.

In conclusion, we have to state that the nature of big difference in baryon and meson production in Au+Au collisions at RHIC in transition region between soft and hard scattering is still not clear, more data are needed, more consistent explanations are required.

References

1. T.K. Hemmick, J. Phys. **G30**, S659 (2004).
2. S.S Adler *et al* (PHENIX Collaboration), *Phys. Rev. Lett.* **91**, 172301 (2003).
3. K. Scheweda (STAR Collaboration), J. Phys. **G30**, S693 (2004).
4. S.S Adler *et al* (PHENIX Collaboration), nucl-ex/0410012, (2004).
5. L.S Barnby (STAR Collaboration), J. Phys. **G30**, S1121 (2004).
6. S. Bathe (PHENIX Collaboration), talk at APS Meeting, October 27-30, Chicago, IL, USA.
7. C. Adler *et al* (STAR Collaboration), *Phys. Rev. Lett.* **90**, 082302 (2003).
8. J. Belchikova (STAR Collaboration), talk at APS Meeting, October 27-30, Chicago, IL, USA.
9. S.S Adler *et al* (PHENIX Collaboration), nucl-ex/0408007, (2004).
10. R.J. Fries *et al*, *Phys. Rev. Lett.* **90**, 202303 (2003). R.C. Hwa and C.B. Yang, *Phys. Rev. C* **70**, 024904 (2004).
11. X. Zhang and G. Fai, hep-ph/0306227 (2004).




Figure 5: Nuclear modification factor at 62.4 GeV for most central and semi-central Au+Au collisions. Top points - for all charged hadrons, lower points - for neutral pions only.




Figure 6: Proton to pion ratio at 62.4 GeV for most central and semi-central Au+Au collisions.





Figure 7: Results from 2-particle correlation with identified leading hadron: yield for near and away side jets per trigger particle vs. number of participants. STAR preliminary results are on the left histograms, PHENIX results - on the right. In PHENIX, baryons were protons and antiprotons, mesons - pions and Kaons; dotted line shows the results of recombination model. Difference in STAR/PHENIX yields is due to about factor 2 larger pseudorapidity acceptance in STAR.